

Consent for Botox for Hyperhidrosis

Botox is indicated for the treatment of severe primary axillary hyperhidrosis (underarm sweating) that is inadequately managed with topical agents.

Please tell us if:

- You are allergic to Botox or anything in the preparation
- You have a neuromuscular disease such as ALS, motor neuropathy myasthenia gravis or Lambert-Eaton syndrome.
- You have an infection at the intended site of injection
- You are taking antibiotics or other prescription medications.

Warnings:

The recommended dosage and frequency of the administration of Botox should not be exceeded.

Serious and/or immediate hypersensitivity reactions have been rarely reported. These reactions include anaphylaxis, hives, soft tissue edema, and shortness of breath.

Individuals with peripheral motor neuropathic diseases or neuromuscular disorders may be at increased risk of clinically significant systemic effects including severe shortness of breath and respiratory compromise from typical doses of Botox.

Adverse Reactions:

The most frequently reported adverse events (3% to 10% of patients) following Botox injection for severe primary axillary hyperhidrosis included injection-site pain and hemorrhage, non-axillary sweating, infection, sore throat, flu syndrome, headache, fever, neck or back pain, itching, and anxiety.

I have read the above information and consent to be treated with Botox for hyperhidrosis.

Signature

Date

Print name

Lot # / Expiration date